



CONTROLS:



Aphids



Cutworms



Fleas



Spider Mites

And other listed insects

VEGETABLES PLUS

INSECTICIDE

- Kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs, vegetables, fruits, and nuts
- Provides contact and residual control for up to 4 weeks
- Kills Over 100 Named Insect Pests
- Multi-Purpose, Broad Spectrum Insecticide

ACTIVE INGREDIENT:

Permethrin.....10.00%

INERT INGREDIENTS:.....90.00%

TOTAL:100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel Booklet for Additional Precautionary Statements and Complete Directions for Use.

A **CONTROL SOLUTIONS INC.** PRODUCT

NET CONTENTS: 1 PINT



0 72693 04881 2



CONTROLS:



Aphids



Cutworms



Fleas



Spider Mites

And other listed
insects

VEGETABLES PLUS

INSECTICIDE

- Kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs, vegetables, fruits, and nuts
- Provides contact and residual control for up to 4 weeks
- Kills Over 100 Named Insect Pests
- Multi-Purpose, Broad Spectrum Insecticide

ACTIVE INGREDIENT:

Permethrin.....10.00%

INERT INGREDIENTS:.....90.00%

TOTAL:.....100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel Booklet for Additional
Precautionary Statements and Complete Directions for Use.

A **CONTROL SOLUTIONS INC.** PRODUCT

NET CONTENTS: 1 QUART





VEGETABLES PLUS INSECTICIDE CONCENTRATE

- Kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs, vegetables, fruits, and nuts
- Provides contact and residual control for up to 4 weeks
- Kills Over 100 Named Insect Pests
- Multi-Purpose, Broad Spectrum Insecticide

ACTIVE INGREDIENT:

Permethrin 10.0%

INERT INGREDIENTS: 90.0%

TOTAL: 100.0%

EPA Reg. No. 53883-69

EPA Est. No. 53883-TX-002

KEEP OUT OF REACH OF CHILDREN CAUTION

READ ENTIRE LABEL AND PAMPHLET.
USE STRICTLY IN ACCORDANCE WITH
PRECAUTIONARY STATEMENTS AND
DIRECTIONS AND WITH APPLICABLE
STATE AND FEDERAL REGULATIONS.

CONTROL SOLUTIONS, INC.
5903 GENOA-RED BLUFF
PASADENA, TEXAS 77507-1041

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

FIRST AID *(continued)*

If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

(continued)

FIRST AID *(continued)*

NOTE TO PHYSICIAN AND VETERINARIANS: Do not administer or instill milk, cream or other substances containing vegetable or animal fats, which will enhance absorption of lipophilic substances.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. Do not apply directly to water. Avoid contamination of ornamental fish ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.
Do not use in undiluted form.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle an open container in a manner as to prevent spillage.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-refillable container. Do not re-use or re-fill this container. Offer for recycling if available. Completely empty container. Dispose of container at an approved waste disposal facility or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed, or drinking water.

Spills: If the container is leaking or material is spilled, absorb liquid with an absorbent such as kitty litter and carefully sweep up and wash exposed area with detergent and water. In spill or leak incidents, keep unauthorized people away.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL PRECAUTIONS AND USE RESTRICTIONS

- Apply this product only as specified on this label.
- Do not use spoons or measuring cups for food purposes after use with pesticides
- Do not use in food areas of restaurants or food handling establishments or other areas where food is commercially prepared or processed. Do not use in a facility that is in operation or in serving areas while food is exposed. Serving areas are defined as areas where prepared foods are served such as dining rooms, but exclude areas where food may be prepared or held. When used in the home, cover or remove all exposed food and cooking utensils and cover all food-processing surfaces or wash thoroughly after treatment and before re-use.
- Do not contaminate food or feedstuffs.

(continued)

GENERAL PRECAUTIONS AND USE RESTRICTIONS *(continued)*

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Do not remain in the treated area. Exit immediately and remain outside the treated area until vapors, mists and aerosols have dispersed.
- Remove or cover exposed food and drinking water before application.
- Do not apply directly to or near water, storm drains or drainage ditches. Do not apply in windy conditions. To prevent run-off, do not water the treated area(s) or apply when heavy rain is expected. Rinse application equipment only over treated area.
- Do not make applications during rain.
- Do not wet plants to point of run-off or drip. Do not use treated articles until spray has dried.
- Remove pets and birds, and cover fish aquaria before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.

(continued)

GENERAL PRECAUTIONS AND USE RESTRICTIONS *(continued)*

- All outdoor applications must be limited to spot or crack-and-crevice applications, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf, and other vegetation; Application to building foundations up to a maximum height of 3 feet. Other than applications to building foundations all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches, and structural surfaces (such as windows, doors and eaves) are limited to crack and crevice applications only.

THREE STEPS TO MIX AND APPLY: SHAKE WELL BEFORE USING

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.
2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.
3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed.

Volume conversions: 1 fl. oz. = 6 tsp. or 2 Tbsp.; 8 fl. oz. = ½ pt. or 1 cup

DIRECTIONS FOR APPLYING

HOME LAWNS

Turf Variety	Pests	How to Use
bentgrass, bermudagrass, bluegrass, dichondra, fescue, Irish moss, merion, St. Augustine grass	ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, fleas, grasshoppers, Japanese beetle grubs, mole crickets, sod webworms, ticks (including those that carry Lyme disease), leafhoppers	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For ants, crickets and grasshoppers, use ½ fl. oz. in 3 gals. of water to cover 1,000 sq. ft. For other listed insects, use 1½ fl. oz. in 10 gal. of water to cover 1,000 sq. ft.

HOUSE PLANTS INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND FOR INDOOR LANDSCAPING:

House Plants	Pests	How to Use
ageratum, ardisia, aster, baby's breath, begonia, birdsnest fern, bleeding heart, carnation, chrysan- themum, coleus, crown of thorns, cockscomb, cyclamen, dracaena, dumbcane, English ivy, fuchsia, gladiolus, grape ivy,	aphids, armyworms, corn earworms, exposed thrips, fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scales, soldier mites, spring cankerworms, whiteflies	Mix ¼ fl. oz. in 1 gal. of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause

(continued)

House Plants	Pests	How to Use
<p><i>(continued)</i> juniper, lily, marigold, orchid, palm, pansy, peperomia, petunia, philodendron, piggy- back plant, poinsettia, portulaca, pothos, prayer plant, purple passion, rabbits foot fern, rose, schefflera, snapdragon, snake plant, statice, velvet plant, verbena, zinnia</p>	<p>aphids, armyworms, corn earworms, exposed thrips, fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scales, soldier mites, spring cankerworms, whiteflies</p>	<p>some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury.</p>
<p>velvet plants</p>	<p>fungus gnats</p>	

**Fire Ant Control
Individual Mound and Plant Container
Treatment:**

Mix ½ fl. oz. in 3 gals. of water and apply as a drench using a sprinkler can. Gently apply 1 gallon of the mixture over the surface of each mound. Thoroughly wet mound and surrounding area of 3x4 ft. (12 sq. ft.). For best results, apply in cool weather 65°-80°F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration, which will reduce product effectiveness. Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.

TREES & SHRUBS

Tree and Shrub Variety	Pests	How to Use
Arizona, cypress, azalea, birch, cherry, non-bearing citrus, conifers, elm, English ivy, euonymus, weeping fig, fir, honeysuckle, lilac, mock-orange, oak, palm, philodendron, pine, poinsettia, tulip poplar, rhododendron, taxus	aphids, bagworms, cicadas, exposed thrips, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, spider mites, whiteflies, gypsy moth, tent caterpillar, Japanese beetles	Use ¼ fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening when the temperature ranges from 50°-75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.

(continued)

Tree and Shrub Variety	Pests	How to Use
cherry, honeysuckle	fall webworms	Use ¼ fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening when the temperature ranges from 50°-75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.
pine	pine beetles, pine moths, pine needle miners, needle scales	
other conifers	budworms	
Douglas and grand firs	tussock moth	
non-bearing citrus	citrus black flies	
oak	oakworms	
elm	elm leaf beetles, elm spanworms, fall webworms	

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

ROSES & FLOWERS (ORNAMENTALS)

Roses & Flowers (Ornamentals)	Pests	How to Use
ageratum, aster, ardisia, azalea, baby's breath, begonia, birds-nest fern, bleeding heart, carnation, chrysanthemum, coleus, common ninebark and snowberry, crown of thorns, cockscomb, cyclamen, dracaena,	aphids, armyworms, bagworms, cabbage worms, cicadas, cornworms, cutworms, exposed thrips, fall cankerworms, fall webworms, fungus gnat, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale crawlers, spring cankerworms, spider mites, whiteflies, rose chafer, psyllids, scale insects	Use ¼ fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.

(continued)

Roses & Flowers (Ornamentals)	Pests	How to Use
dumbcane, English ivy, exacum, fuchsia, gladiolus, grape ivy, gold bells, hypoestes, ivy, juniper, lilac, marigold, mock-orange, nannyberry, orchid, palm, pansy, pea shrub, peperomia, petunia, philodendron, piggy-back plant,	aphids, armyworms, bagworms, cabbage worms, cicadas, cornworms, cutworms, exposed thrips, fall cankerworms, fall webworms, fungus gnat, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale crawlers, spring cankerworms, spider mites, whiteflies, rose chafer, psyllids, scale insects	Use ¼ fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.

(continued)

Roses & Flowers (Ornamentals)	Pests	How to Use
<p>poinsettia, portulaca, pothos, prayer plant, purple passion, rabbits foot fern, rhododendron, rose, schefflera, snapdragon, snake plant, statice velvet plant, verbena, zinnia</p>	<p>aphids, armyworms, bagworms, cabbage worms, cicadas, cornworms, cutworms, exposed thrips, fall cankerworms, fall webworms, fungus gnat, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale crawlers, spring cankerworms, spider mites, whiteflies, rose chafer, psyllids, scale insects</p>	<p>Use ¼ fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.</p>

FRUIT & NUT TREES
GENERAL DIRECTIONS

Use ½ fl. oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Cover undersides of leaves and penetrate dense foliage to the extent possible.

Tree Variety	Pests	How to Use
almonds, pecans	navel orangeworms, peach twig borers	Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.

(continued)

Tree Variety	Pests	How to Use
apples	green fruitworms, aphids, leafrollers (oblique-banded, redbanded), plum curculio, rosy apple aphids, spotted tentiform leafminers, tarnished plant bugs, white apple leafhoppers, Japanese beetles	Do not make more than 3 applications. Do not apply after petal fall.
peaches	borers (lesser peachtree, peach twig), green fruitworms, plum curculio, Oriental fruit moths, tarnished plant bugs	Do not apply within 7 days of harvest. Do not make more than 8 applications per season.

(continued)

Tree Variety	Pests	How to Use
pears (dormant through delayed dormant)	pear psylla	Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.
pears (summer)	codling moths, green fruitworms, pear psylla, aphids	Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

VEGETABLES

Prepare a spray solution at the dilution rates listed for each crop in the chart below. Apply the spray solution at the rate of 1 gallon to cover 440 square feet surface area.

Vegetables	Pests	How to Use
asparagus (pre-harvest)	asparagus beetles, cutworms	Use $\frac{3}{4}$ fl. oz. in 2 gals. of water. Apply to ensure uniform coverage. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
asparagus (post-harvest)	larval stages of asparagus beetle; tarnished plant bugs, lygus bugs, adult Japanese beetle	Use $\frac{1}{4}$ fl. oz. in 1 gal. of water. Treat ferns or bush growth after spear harvest when insects are present.

(continued)

Vegetables	Pests	How to Use
<i>celery</i> - see leafy vegetables		
cole crops, broccoli, Brussels sprouts, cabbage, cauliflower	beet armyworms, cabbage aphids and loopers, diamondback moths, imported cabbageworms	Use $\frac{1}{4}$ fl. oz. in 1 gal. of water. Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on cauliflower, Brussels sprouts and broccoli and 10 applications on cabbage.
<i>corn</i> - see sweet corn		
cucumber, melons, pumpkins, squash	flea beetle, leafhopper	Use $\frac{1}{2}$ fl. oz. in 1 gal. water. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.

(continued)

Vegetables	Pests	How to Use
<u>cucurbits</u> : balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey	aphids, cabbage looper, cucumber beetle (adults), cutworms, leafhoppers, leafminers, melonworms, pickleworm, plant bugs, squash bugs	Use ½ fl. oz. in 1 gal. water. Apply to ensure uniform coverage. Can be applied up to the day of harvest.

(continued)

Vegetables	Pests	How to Use
<i>(continued)</i> balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon	aphids, cabbage looper, cucumber beetle (adults), cutworms, leafhoppers, leafminers, melonworms, pickleworm, plant bugs, squash bugs	Use ½ fl. oz. in 1 gal. water. Apply to ensure uniform coverage. Can be applied up to the day of harvest.
eggplant	Colorado potato beetles, vegetable leafminers	Use ¾ fl. oz. in 1 gal. of water. Apply to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.

(continued)

Vegetables	Pests	How to Use
horseradish (foliar application)	imported crucifer weevils	Use $\frac{3}{4}$ fl. oz. in 1 gal. water. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.
head lettuce	corn earworms, beet armyworms, loopers (alfalfa, cabbage), tobacco budworms	Use $\frac{1}{2}$ fl. oz. in 1 gal. water. Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.

(continued)

Vegetables	Pests	How to Use
<p><u>leafy vegetables:</u> amaranth, arrugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible-leaved and garland), cress (garden and upland), dandelion, dock, endive, fennel, lettuce (head and leaf), okra, parsley, purslane (garden and winter), rhubarb, spinach, Swiss chard</p>	<p>alfalfa looper, aphids, beet armyworm, cabbage looper, corn earworm, cutworms, european corn borer, fall armyworm, green cloverworm, southern armyworm, tobacco budworm, vegetable leafminer</p>	<p>Use $\frac{1}{3}$ fl. oz. in 1 gal. water. Apply every 5 to 10 days as needed. Can be applied up to 1 day prior to harvest.</p>

(continued)

Vegetables	Pests	How to Use
melons - see cucurbits		
potatoes	beet armyworm, Colorado potato beetle, potato leafhopper, potato tuberworm, cabbage looper, cutworms, European corn borer, potato psyllid, tarnished plant bug	Use $\frac{3}{4}$ fl. oz. in 2 gals. of water. Apply as needed, but not more than 12 applications. Can be applied up to 7 days prior to harvest.

(continued)

Vegetables	Pests	How to Use
peppers	vegetable leafminers, cabbage looper, flea beetle, corn earworm, pepper weevil, European corn borer, green cloverworms, leafhoppers	Use ½ fl. oz. in 1 gal. of water. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.
strawberries, raspberries, blueberries	strawberry weevils, aphids, spider mites, lygus bugs, flea beetles, omnivorous leaf-tier, spittlebugs, Japanese beetles, field crickets, red spider, leafroller, weevils	Use ½ fl. oz. in 1 gal. water. Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.

(continued)

Vegetables	Pests	How to Use
sweet corn	armyworms (fall, southern), corn earworms, european corn borers	Use $\frac{3}{4}$ fl. oz. in 2 gals. of water. Apply every 5 days, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
tomatoes	beet and southern armyworm, tomato pinworm, tomato fruitworm, horn worms, granulate cutworms, Colorado potato beetle, cabbage looper	Use $\frac{1}{2}$ fl. oz. in 1 gal. water. Apply to ensure uniform coverage. Can be applied up to day of harvest.

NUISANCE PESTS IN OUTDOOR AREAS

GENERAL DIRECTIONS

Use 1 $\frac{2}{3}$ fl. ozs. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.

OUTSIDE SURFACES OF BUILDING

Use Vegetables Plus on ants, clover mites, crickets, earwigs, millipedes, sowbugs (pillbugs) and spiders to help prevent infestation of buildings. Treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen.

OTHER OUTDOOR AREAS

Use Vegetables Plus on ants, crickets, earwigs, fleas, ticks (**including ticks that may carry Lyme disease**). For treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. **For ants**, thoroughly wet hills and runways. Repeat application only as needed to maintain insect control.

To reduce annoyance from: **cluster flies, flying moths, gnats, house flies, mosquitoes.** Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate.

ANIMALS AND ANIMAL PREMISE USE

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 ounces per gallon) (0.5%). For normal infestations, dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon of water) ($1\frac{1}{3}$ tablespoonful per gallon) (0.1%).

PESTS ON DOG PREMISES
(Kennels, Dog Houses, Runs and Yards)

33

Fleas, Flies, Ticks, Lice and Mange Mites (adult and larvae)	Dilute 8 oz. in 6.25 gal. water Dilute 1 pt. in 12.5 gal. water Dilute 1 qt. in 25 gal. water	Thoroughly spray infested areas, pet beds, and resting quarters. Fresh bedding should be placed in animal quarters following treatment. Most efficient pest control is accomplished when heavy pressure and good contact is achieved. Timing and frequency of applications should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray food, feed, or water. Repeat as needed but not more than once every two weeks.
--	---	--

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.

WARRANTY STATEMENT

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent consistent with applicable law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency.

Except, as expressly provided herein, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.

NOTES



VEGETABLES PLUS

INSECTICIDE CONCENTRATE

- Kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs, vegetables, fruits, and nuts
- Provides contact and residual control for up to 4 weeks
- Kills Over 100 Named Insect Pests
- Multi-Purpose, Broad Spectrum Insecticide

ACTIVE INGREDIENT:

Permethrin 10.0%

INERT INGREDIENTS: 90.0%

TOTAL: 100.0%

EPA Reg. No. 53883-69

EPA Est. No. 53883-TX-002

KEEP OUT OF REACH OF CHILDREN CAUTION

READ ENTIRE LABEL AND PAMPHLET.
USE STRICTLY IN ACCORDANCE WITH
PRECAUTIONARY STATEMENTS AND
DIRECTIONS AND WITH APPLICABLE
STATE AND FEDERAL REGULATIONS.

CONTROL SOLUTIONS, INC.
5903 GENOA-RED BLUFF
PASADENA, TEXAS 77507-1041

0711