

UHS

Team* 2G

Herbicide

A selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bahiagrass, Bermudagrass, Bluegrass (perennial), Centipedegrass, Fescue, Ryegrass (perennial), St. Augustinegrass, Zoysiagrass, Bentgrass Fairways, and other high cut Bentgrass areas

ACTIVE INGREDIENTS:

benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine 1.33%
trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.67%

INERT INGREDIENTS: 98.00%

TOTAL 100.00%

Contains 1 pound active ingredient per 50 pound bag

*TEAM and the TEAM logo are trademarks of Dow AgroSciences, LLC

EPA REG. NO. 62719-137-65783

EPA EST NO. 961-PA^(L)-01:961-IL^(D)-01

Superscript used is first letter of lot number.

IHT

01/00

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

See Below For Additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION: Causes eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with skin, eyes, or clothing. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation develops.

If inhaled: If discomfort occurs, remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and call a physician.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" elsewhere on this label.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

TEAM 2G HERBICIDE is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and

hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia supina*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*). TEAM 2G HERBICIDE controls susceptible annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds.

When TEAM 2G HERBICIDE is applied as directed, the following established turfgrasses are tolerant:

Bahiagrass	St. Augustinegrass
Bermudagrass	Zoysiagrass
Centipedegrass	Bentgrass Fairways,
Fescue	and other high cut
Kentucky bluegrass	Bentgrass areas
Perennial ryegrass	

TEAM 2G HERBICIDE will not harm most nearby established ornamental trees and shrubs.

Do not apply this product to sod farms, turfgrasses grown for seed, farms, greenhouses or nurseries, including plants or plant materials grown for sale or research purposes.

General Use Precautions

Apply TEAM 2G HERBICIDE to the turf prior to the germination of weed grasses since this product will not control established weeds. TEAM 2G HERBICIDE may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 100 pounds per acre (2 pounds active ingredient) or 2 1/4 pounds per 1000 square feet. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been over-seeded with winter grasses, a spring application of TEAM 2G HERBICIDE may thin the over-seeded grasses. Because of varying cultural practices, this product should not be used on putting greens. TEAM 2G HERBICIDE should not be used on dichondra. TEAM 2G HERBICIDE should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until these turfs are well established.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply TEAM 2G HERBICIDE approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance should be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application of TEAM 2G HERBICIDE.

1. Spring Germinating Summer Annual Grasses—Crabgrasses, Goosegrass, Foxtails and Barnyardgrass

A. Northern (Cool Season) Turfgrasses

For preemergence control, apply 75 to 150 pounds per acre (1.5 to 3 pounds active ingredient) or 1 3/4 to 3 1/2 pounds per 1000 square feet of TEAM 2G HERBICIDE in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination. Where heavy goosegrass pressure exists, use the rate of 150 pounds of this product per acre or 3 1/2 pounds per 1000 square feet. If split applications are desired, apply a cumulative rate of 150 pounds per acre (3 pounds active ingredient) or 3 1/2 pounds per 1000 square feet and make the second application 8 to 10 weeks after the first.

B. Southern (Warm Season) Turfgrasses

For preemergence control, apply 100 to 150 pounds per acre (2 to 3 pounds active ingredient) or 2 1/4 to 3 1/2 pounds per 1000 square feet of TEAM 2G HERBICIDE in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of TEAM 2G HERBICIDE at the recommended rate may be made 10 to 12 weeks after initial treatment for continued control.

2. *Poa annua* (Annual Bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in thin areas previously occupied by annual bluegrass.

A. Northern (Cool Season) Turfgrasses

For preemergence control, apply 75 to 100 pounds per acre (1 1/2 to 2 pounds active ingredient) or 1 3/4 to 2 1/4 pounds per 1000 square feet of TEAM 2G HERBICIDE in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

B. Southern (Warm Season) Turfgrasses

For preemergence control, apply 150 pounds per acre (3 pounds active ingredient) or 3 1/2 pounds per 1000 square feet of TEAM 2G HERBICIDE in the late summer or early fall one to two weeks prior to the expected

germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application of TEAM 2G HERBICIDE at 150 pounds per acre can be made 10 to 12 weeks after the initial treatment. For application to smaller areas, apply the following amounts of TEAM 2G HERBICIDE.

TEAM 2G HERBICIDE/Acre	TEAM 2G HERBICIDE/1000 Sq Ft
75 lbs.	1 ¼ lbs.
100 lbs.	2 ¼ lbs.
150 lbs.	3 ½ lbs.

reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. Reseeding should be delayed 8 weeks after application of TEAM 2G HERBICIDE at the 75 to 100 pounds per acre (1.5 to 2 pounds active ingredient) or 1 ¼ to 2 ¼ pounds per 1000 square feet rate and 12-16 weeks at the 100 to 150 pounds per acre (2 to 3 pounds active ingredient) or 2 ¼ to 3 ½ pounds per 1000 square feet rate.


APPLICATION EQUIPMENT

Apply TEAM 2G HERBICIDE evenly over the turfgrass area without skips or misses. A more uniform application can be made by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. For best results, use equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory

Helpful Application Techniques For Applying TEAM 2G HERBICIDE



Appropriate Spreader Settings As A Guide For Calibration (for applying TEAM 2G HERBICIDE in one pass over the turf)

Note: These settings are provided as suggested starting points in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by United Horticultural Supply or the spreader manufacturer.

Spreader Type	Spreader Setting Necessary To Apply Pounds Formulated Product Per Acre		
	75 lbs./Acre	100 lbs./Acre	150 lbs./Acre
Vicon: Model 402 (with seed ring) Swath width 20 ft, 5 mph speed	19	24	28

Spreader Type	Spreader Setting Necessary To Apply Pounds Formulated Product Per Acre		
	75 lbs./Acre	100 lbs./Acre	150 lbs./Acre
Gandy: Model 10A series, 6, 8, 10, 11, 12, 20 and 30 ft. (based on tractor speed of 5 mph)	26	30	36
Model 10T series, 6, 8, 10, 11, 12, 20, and 30 ft. (based on tractor speed of 5 mph)	28	32	39
Turf Tender, 24, 30, 42 inch (based on tractor speed of 3 mph)	24	26	28.5
600 Series, 2, 4, and 5 ft. (based on tractor speed of 3 mph)	15.5	18	22
Scott: Rotary Model R-8	G	G ½	1 ½
Drop Spreader	3 ½	3 ¾	4 5/8
Lesco: Lesco Spreader (Note: set 3rd adjustable hole 1 notch from 'open')	F	F ½	H ¼
Cyclone: Models B1, 33, 96, BSSB and BPI at 200 ft/min.	3 1/8	3 ¼	3 ½
Model 100 at 200 ft/min.	3 ¼	3 ½	3 ¾
Model 93 (based on tractor speed of 4 mph)	4 ½	5 ¼	6 ½
Model 99 (based on tractor speed of 4 mph)	4 ¼-4 ½	4 ¾-5	6
Model BPI (based on tractor speed of 4 mph)	4-4 ¼	4 ¾	5 ½
Spyker: Spyker	3.1	3.3	3.7
Lely: Model W and H (based on tractor speed of 4.5 mph)	2	3	4

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs or Clothing.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

FORMULATED FOR UNITED HORTICULTURAL SUPPLY